



Sandiway Primary School First Aid and Medication Policy

June 2022



Sandiway Primary School is committed to caring for, and protecting, the health, safety and welfare of its pupils, staff and visitors. We confirm our adherence to the following standards at all times:

- To make practical arrangements for the provision of First Aid on our premises, during off-site sport and on school visits.
- To ensure that trained First Aid staff renew, update or extend their HSE approved qualifications at least every three years.
- To have the minimum of trained First Aiders on site at any one time, including a person with a paediatric first aid qualification whenever EYFS pupils are present. Such people will be able to responsibly deliver or organise emergency treatment.
- To ensure that a trained first aider accompanies every off-site visit and activity. In visits involving EYFS pupils, such a person will have a current paediatric first aid qualification.
- To record accidents and illnesses appropriately, reporting to parents and the Health & Safety Executive under the Reporting of Injuries, Diseases and Dangerous Occurrences Regulations (1995).
- To provide accessible first aid kits at various locations on site, along with a portable kit for trips, excursions and sport.
- To record and make arrangements for pupils and staff with specific medical conditions.
- To deal with the disposal of bodily fluids and other medical waste accordingly, providing facilities for the hygienic and safe practice of first aid.
- To contact the medical emergency services if they are needed, informing next of kin immediately in such a situation.
- To communicate clearly to pupils and staff where they can find medical assistance if a person is ill or an accident has occurred.
- To communicate clearly in writing to parents or carers if a child has sustained a bump to the head at school, however minor, and to communicate in writing in relation to every instance of accident or first aid or the administration of medicine for pupils in EYFS.

Details of First Aid Practitioners at Sandiway Primary School

Found displayed in school

Practical Arrangements at Sandiway Primary School

Location of First Aid Facilities

- The sick area is located next to the staff room for first aid treatment and for pupils or staff to rest/recover if feeling unwell. This includes: first aid supplies, a water supply and sink, an adjacent bathroom and hygiene supplies such as gloves and paper towels.
- An additional first aid kit is also available in this area for quick release off a bracket to take to an incident.
- The defibrillator is also in this area of the school
- The music room is available as a more private space or if a child/adult needs to be isolated in the event of a suspected COVID-19 case.

A portable first aid kit must be obtained from the office for school visits.

Responsibilities of the Trained First Aiders

- Provide appropriate care for pupils or staff who are ill or sustain an injury
- Record all accidents in the accident book (to be found next to the staff room). They are then passed to the school secretary who will make a copy for individual pupil files.
- In the event of any injury to the head, however minor, ensure that a note from the office is sent home to parents/carers and a copy placed in the pupil's file. A fully qualified first-aider is responsible for this part of the treatment.
- In the event of any accident or administration of first aid involving a pupil in EYFS, ensure that a note from the office is sent home to parents/carers.
- Make arrangements with parents/carers to collect children and take them home if they are deemed too unwell to continue the school day.
- Inform the appointed person of all incidents where first aid has been administered.

Responsibilities of the Appointed Person

- Ensure that all staff and pupils are familiar with the school's first aid and medical procedures.
- Ensure that all staff are familiar with measures to provide appropriate care for pupils with particular medical needs (eg. diabetic needs, Epi-pens, inhalers).
- Blue medical boxes are kept in the office out of reach of the children, all staff are given code for the box, any medication requiring refrigeration will be kept in the kitchen fridge, kitchen is always kept locked.
- Ensure that a list is maintained and available to staff of all pupils with particular medical needs and appropriate measures needed to care for them.
- Monitor and re-stock supplies and ensure that first aid kits are replenished.
- Ensure that the school has an adequate number of appropriately trained First Aiders.
- Co-ordinate First Aiders and arrange for training to be renewed as necessary.
- Maintain adequate facilities.
- Ensure that correct provision is made for pupils with special medical requirements both in school and on off-site visits.
- On a regular basis, review First Aid records to identify any trends or patterns and report to the Health and Safety committee
- Fulfil the school's commitment to report to RIDDOR, as described below
- Liaise with managers of external facilities, such as the local sports facilities, to ensure appropriate first aid provision.
- Contact emergency medical services as required.
- Maintain an up-to-date knowledge and understanding of guidance and advice from appropriate agencies

What to do in the case of an accident, injury or illness

- A member of staff or pupil witnessing an accident, injury or illness should immediately contact a named trained first aider (see above).
- The school office should be contacted if the location of a trained first aider is uncertain. Any pupil or member of staff sustaining an injury whilst at school should be seen by a first aider who will provide immediate first aid and summon additional help as needed.
- The pupil or member of staff should not be left unattended.
- The first aider will organise an injured pupil's transfer to the sick room if possible and appropriate and to hospital in the case of an emergency.
- Parents should be informed as necessary by telephone by the first aider or school secretary.
- This will be followed up in writing and a record kept at school. A written record of all accidents and injuries is maintained in the accident book.

Contacting parents

- Parents should be informed by telephone as soon as possible after an emergency or following a serious/significant injury including:
 - Head injury (a head injury advice sheet should be given to any pupil who sustains a head injury) Available from the appointed person
 - Suspected sprain or fracture
 - Following a fall from height
 - Dental injury
 - Anaphylaxis & following the administration of an Epi-pen
 - Epileptic seizure
 - Severe hypoglycaemia for pupils, staff or visitors with diabetes
 - Severe asthma attack
 - Difficulty breathing
 - Bleeding injury (uncontrolled)
 - Loss of consciousness
 - If the pupil is generally unwell

If non-emergency transportation is required, an authorised taxi service or staff car will be used if parents are delayed. A member of staff will accompany the pupil until a parent arrives. Parents can be informed of smaller incidents at the end of the school day by the form teacher.

Contacting the Emergency Services

An ambulance should be called for any condition listed above or for any injury that requires emergency treatment. Any pupil taken to hospital by ambulance must be accompanied by a member of staff until a parent arrives. All cases of a pupil becoming unconsciousness (not including a faint) or following the administration of an Epi-pen, must be taken to hospital.

Accident reporting

- The accident book must be completed for any accident or injury occurring at school, at the local sports facilities, or on a school trip. This includes any accident involving staff or visitors. The accident book will be monitored by the appointed person as certain injuries require reporting (RIDDOR requirements).
- Any pupil who is unwell cannot be left to rest unsupervised in the sick room. If a pupil becomes unwell, a parent should be contacted as soon as possible by the appointed person, the school secretary or the head teacher.
- Anyone not well enough to be in school should be collected as soon as possible by a parent. Staff should ensure that a pupil who goes home ill remembers to sign out at the school office. Informal diagnostic report of child is made, this is kept in the office.

First Aid equipment and materials

The appointed person is responsible for stocking and checking the first aid kits. Staff are asked to notify the appointed person when supplies have been used in order that they can be restocked. The first aid boxes contain:

- A first aid guidance card
- Adhesive hypo allergenic plasters
- Triangular bandages (slings)
- Safety pins
- Cleaning wipes
- Adhesive tape
- Sterile eye pads
- Medium sized unmedicated dressings

- Large sized unmedicated dressings
- Disposable gloves
- Resuscitator
- Yellow clinical waste bag

First aid for school trips

- The trip organiser must ensure that at least one adult accompanying the trip has an appropriate first aid qualification (paediatric certificate for trips involving EYFS pupils) and undertake a risk assessment to ensure an appropriate level of first aid cover, with reference to the educational visits policy, which includes further guidance.
- A First Aid kit for school trips must be collected from the main office (Registrar). This must be returned to the office for replenishing on return.
- Any accidents/injuries must be reported to the appointed person and to parents and documented in the accident book in accordance with this policy.
- RIDDOR guidelines for reporting accidents must be adhered to.
- For any major accident or injury the appropriate health & safety procedure must be followed.

Pupils using crutches or having limited mobility

Parents must inform the school of the nature of injury and the anticipated duration of immobility. The form tutor will arrange for a 'class partner' to carry books, open doors etc. Information about the condition will be discussed in staff meetings to enable teachers to be fully aware of the pupil's needs. Arrangements will be made for the pupil to arrive/leave lessons early to allow for a safe transfer around school. Parents must inform the school of any particular difficulties.

Emergency care plans and treatment boxes

The appointed person ensures that staff are made aware of any pupil with an emergency care plan. These care plans are displayed in the staff room. Pupils with a serious medical condition will have an emergency care plan drawn up and agreed by the appointed person and parents. Emergency treatment boxes must always be taken if the pupil is out of school.

Pupils with medical conditions

A list is available in the staff room of all pupils who have a serious allergy or medical condition. This information is useful for lesson planning and for risk assessments prior to a school trip. Please return emergency boxes on completion of the trip. If staff become aware of any condition not on these lists please inform the appointed person.

Dealing with body fluids

In order to maintain protection from disease, all body fluids should be considered infected. To prevent contact with body fluids the following guidelines should be followed.

- When dealing with any body fluids wear disposable gloves.
- Wash hands thoroughly with soap and warm water after the incident.
- Keep any abrasions covered with a plaster.
- Spills of the following body fluids must be cleaned up immediately.

Bodily fluids include: Blood, Faeces, Nasal and eye discharges, Saliva, Vomit

- Disposable towels should be used to soak up the excess, and then the area should be treated with a disinfectant solution.
- Never use a mop for cleaning up blood and body fluid spillages. All contaminated material should be disposed of in a yellow clinical waste bag (available in all 1st aid boxes) then placed in the outdoor waste bin.

- Avoid getting any body fluids in your eyes, nose, mouth or on any open sores. If a splash occurs, wash the area well with soap and water or irrigate with copious amounts of saline.

Infectious diseases

If a child is suspected of having an infectious disease advice should be sought from the appointed person who will follow the Health Protection Agency guidelines below to reduce the transmission of infectious diseases to other pupils and staff. Further details in Appendix 2

ILLNESS	PERIOD OF EXCLUSION	COMMENTS
COVID-19	10 days from symptoms, 10 days if in contact with someone either close friends/families or informed by NHS test and trace	Symptoms: High temperature and persistent coughing, loss of taste and smell. https://www.nhs.uk/conditions/coronavirus-covid-19/symptoms/
Chickenpox	5 days from onset of rash	Pregnant women up to 20 weeks and those in last 3 weeks of pregnancy should inform their midwife that they have been in contact with chickenpox. Any children being treated for cancer or on high doses of steroids should also seek medical advice.
German Measles	For 5 days from onset of rash	Pregnant women should inform their midwife about contact
Impetigo	Until lesions are crusted or healed	Antibiotic treatment by mouth may speed healing
Measles	5 days from onset of rash	Any children being treated for cancer or on high doses of steroids must seek medical advice
Scabies	Until treatment has been commenced	Two treatments one week apart for cases. Treatment should include all household members and any other very close contacts
Scarlet Fever	5 days after commencing antibiotics	Antibiotic treatment recommended
Slapped Cheek Syndrome	None	Pregnant women up to 20 weeks must inform their midwife about contact
Diarrhoea and vomiting	48 hours from last episode of diarrhoea or vomiting	Exclusion from swimming may be needed

Hepatitis A	Exclusion may be necessary	Consult the Health Protection Agency
Meningococcal meningitis	Until recovered	Communicable disease control will give advice on any treatment needed and identify contact requiring treatment. No need to exclude siblings or other close contacts.
Viral Meningitis	Until fully recovered	Milder illness
Threadworms	None	Treatment is recommended for the pupil and family members
Mumps	5 days from onset of swollen glands	
Head Lice	None	Treatment is recommended for the pupil and close contacts if live lice are found
Conjunctivitis	None	Children do not usually need to stay off school with conjunctivitis if they are feeling well. If, however, they are feeling unwell with conjunctivitis they should stay off school until they feel better
Influenza	Until fully recovered	
Cold sores	None	Avoid contact with the sores
Warts, verrucae	None	Verrucae should be covered in swimming pools, gymnasiums and changing rooms
Glandular fever	None	
Tonsillitis	None	

Appendix 1:

https://www.publichealth.hscni.net/sites/default/files/Guidance_on_infection_control_in%20schools_poster.pdf

Guidance on infection control in schools and other childcare settings



March 2017

Prevent the spread of infections by ensuring: routine immunisation, high standards of personal hygiene and practice, particularly handwashing, and maintaining a clean environment. Please contact the Public Health Agency **Health Protection Duty Room (Duty Room) on 0300 555 0119** or

visit www.publichealth.hscni.net or www.gov.uk/government/organisations/Public-health-england if you would like any further advice or information, including the latest guidance. Children with rashes should be considered infectious and assessed by their doctor.

Rashes and skin infections	Recommended period to be kept away from school, nursery or childminders	Comments
Athlete's foot	None	Athlete's foot is not a serious condition. Treatment is recommended
Chickenpox*	Until all vesicles have crusted over	See: Vulnerable children and female staff – pregnancy
Cold sores, (Herpes simplex)	None	Avoid kissing and contact with the sores. Cold sores are generally mild and self-limiting
German measles (rubella)*	Four days from onset of rash (as per "Green Book")	Preventable by immunisation (MMR x2 doses). See: Female staff – pregnancy
Hand, foot and mouth	None	Contact the Duty Room if a large number of children are affected. Exclusion may be considered in some circumstances
Impetigo	Until lesions are crusted and healed, or 48 hours after commencing antibiotic treatment	Antibiotic treatment speeds healing and reduces the infectious period
Measles*	Four days from onset of rash	Preventable by vaccination (MMR x2). See: Vulnerable children and female staff – pregnancy
Molluscum contagiosum	None	A self-limiting condition
Ringworm	Exclusion not usually required	Treatment is required
Roseola (infantum)	None	None
Scabies	Child can return after first treatment	Household and close contacts require treatment
Scarlet fever*	Child can return 24 hours after commencing appropriate antibiotic treatment	Antibiotic treatment recommended for the affected child. If more than one child has scarlet fever contact PHA Duty Room for further advice
Slapped cheek (fifth disease or parvovirus B19)	None once rash has developed	See: Vulnerable children and female staff – pregnancy
Shingles	Exclude only if rash is weeping and cannot be covered	Can cause chickenpox in those who are not immune i.e. have not had chickenpox. It is spread by very close contact and touch. If further information is required, contact the Duty Room. See: Vulnerable Children and Female Staff – Pregnancy
Warts and verrucae	None	Verrucae should be covered in swimming pools, gymnasiums and changing rooms

Diarrhoea and vomiting illness	Recommended period to be kept away from school, nursery or childminders	Comments
Diarrhoea and/or vomiting	48 hours from last episode of diarrhoea or vomiting	
<i>E. coli</i> O157 VTEC*	Should be excluded for 48 hours from the last episode of diarrhoea	Further exclusion is required for young children under five and those who have difficulty in adhering to hygiene practices
Typhoid* [and paratyphoid] (enteric fever)	Further exclusion may be required for some children until they are no longer excreting	Children in these categories should be excluded until there is evidence of microbiological clearance. This guidance may also apply to some contacts of cases who may require microbiological clearance
Shigella* (dysentery)		Please consult the Duty Room for further advice
Cryptosporidiosis*	Exclude for 48 hours from the last episode of diarrhoea	Exclusion from swimming is advisable for two weeks after the diarrhoea has settled

Respiratory infections	Recommended period to be kept away from school, nursery or childminders	Comments
Flu (influenza)	Until recovered	See: Vulnerable children
Tuberculosis*	Always consult the Duty Room	Requires prolonged close contact for spread
Whooping cough* (pertussis)	48 hours from commencing antibiotic treatment, or 21 days from onset of illness if no antibiotic treatment	Preventable by vaccination. After treatment, non-infectious coughing may continue for many weeks. The Duty Room will organise any contact tracing necessary

Other infections	Recommended period to be kept away from school, nursery or childminders	Comments
Conjunctivitis	None	If an outbreak/cluster occurs, consult the Duty Room
Diphtheria*	Exclusion is essential. Always consult with the Duty Room	Family contacts must be excluded until cleared to return by the Duty Room. Preventable by vaccination. The Duty Room will organise any contact tracing necessary
Glandular fever	None	
Head lice	None	Treatment is recommended only in cases where live lice have been seen
Hepatitis A*	Exclude until seven days after onset of jaundice (or seven days after symptom onset if no jaundice)	The duty room will advise on any vaccination or other control measure that are needed for close contacts of a single case of hepatitis A and for suspected outbreaks
Hepatitis B*, C, HIV/AIDS	None	Hepatitis B and C and HIV are bloodborne viruses that are not infectious through casual contact. For cleaning of body fluid spills. See: Good Hygiene Practice
Meningococcal meningitis*/septicaemia*	Until recovered	Some forms of meningococcal disease are preventable by vaccination (see immunisation schedule). There is no reason to exclude siblings or other close contacts of a case. In case of an outbreak, it may be necessary to provide antibiotics with or without meningococcal vaccination to close contacts. The Duty Room will advise on any action needed.
Meningitis* due to other bacteria	Until recovered	Hib and pneumococcal meningitis are preventable by vaccination. There is no reason to exclude siblings or other close contacts of a case. The Duty Room will give advice on any action needed
Meningitis viral*	None	Milder illness. There is no reason to exclude siblings and other close contacts of a case. Contact tracing is not required
MRSA	None	Good hygiene, in particular handwashing and environmental cleaning are important to minimise any danger of spread. If further information is required, contact the Duty Room
Mumps*	Exclude child for five days after onset of swelling	Preventable by vaccination (MMR x2 doses)
Threadworms	None	Treatment is recommended for the child and household contacts
Tonsillitis	None	There are many causes, but most cases are due to viruses and do not need an antibiotic

* denotes a notifiable disease. It is a statutory requirement that doctors report a notifiable disease to the Director of Public Health via the Duty Room.

Outbreaks: if a school, nursery or childminder suspects an outbreak of infectious disease, they should inform the Duty Room.

Good hygiene practice

Handwashing is one of the most important ways of controlling the spread of infections, especially those that cause diarrhoea and vomiting, and respiratory disease. The recommended method is the use of liquid soap, warm water and paper towels. Always wash hands after using the toilet, before eating or handling food, and after handling animals. Cover all cuts and abrasions with waterproof dressings.

Coughing and sneezing easily spread infections. Children and adults should be encouraged to cover their mouth and nose with a tissue. Wash hands after using or disposing of tissues. Spitting should be discouraged.

Personal protective equipment (PPE). Disposable non-powdered vinyl or latex-free CE-marked gloves and disposable plastic aprons must be worn where there is a risk of splashing or contamination with blood/body fluids (for example, nappy or pad changing). Goggles should also be available for use if there is a risk of splashing to the face. Correct PPE should be used when handling cleaning chemicals.

Cleaning of the environment, including toys and equipment, should be frequent, thorough and follow national guidance. For example, use colour-coded equipment, follow Control of Substances Hazardous to Health (COSHH) regulations and correct decontamination of cleaning equipment. Monitor cleaning contracts and ensure cleaners are appropriately trained with access to PPE.

Cleaning of blood and body fluid spillages. All spillages of blood, faeces, saliva, vomit, nasal and eye discharges should be cleaned up immediately (always wear PPE). When spillages occur, clean using a product that contains both a detergent and a disinfectant. Use as per manufacturer's instructions and ensure it is effective against bacteria and viruses and suitable for use on the affected surface. Never use mops for cleaning up blood and body fluid spillages – use disposable paper towels and discard clinical waste as described below. A spillage kit should be available for blood spills.

Laundry should be dealt with in a separate dedicated facility. Soiled linen should be washed separately at the hottest wash the fabric will tolerate. Wear PPE when handling soiled linen. Children's soiled clothing should be bagged to go home, never rinsed by hand.

Clinical waste. Always segregate domestic and clinical waste, in accordance with local policy. Use nappies/pads, gloves, aprons and soiled dressings should be stored in correct clinical waste bags in foot-operated bins. All clinical waste must be removed by a registered waste contractor. All clinical waste bags should be less than two-thirds full and stored in a dedicated, secure area while awaiting collection.

Sharps, eg needles, should be discarded straight into a sharps bin conforming to BS 7320 and UN 3291 standards. Sharps bins must be kept off the floor (preferably wall-mounted) and out of reach of children.

Sharps injuries and bites

If skin is broken as a result of a used needle injury or bite, encourage the wound to bleed/wash thoroughly using soap and water. Contact GP or occupational health or go to A&E immediately. Ensure local policy is in place for staff to follow. Contact the Duty Room for advice, if unsure.

Animals

Animals may carry infections, so wash hands after handling animals. Health and Safety Executive for Northern Ireland (HSENI) guidelines for protecting the health and safety of children should be followed.

Animals in school (permanent or visiting). Ensure animals' living quarters are kept clean and away from food areas. Waste should be disposed of regularly, and litter boxes not accessible to children. Children should not play with animals unsupervised. Hand-hygiene should be supervised after contact with animals and the area where visiting animals have been kept should be thoroughly cleaned after use. Veterinary advice should be sought on animal welfare and animal health issues and the suitability of the animal as a pet. Reptiles are not suitable as pets in schools and nurseries, as all species carry salmonella.

Visits to farms. For more information see <https://www.hseni.gov.uk/publications/preventing-or-controlling-ill-health-animal-contact-visitor-attractions>

Vulnerable children

Some medical conditions make children vulnerable to infections that would rarely be serious in most children, these include those being treated for leukaemia or other cancers, on high doses of steroids and with conditions that seriously reduce immunity. Schools and nurseries and childminders will normally have been made aware of such children. These children are particularly vulnerable to chickenpox, measles and parvovirus B19 and, if exposed to either of these, the parent/carer should be informed promptly and further medical advice sought. It may be advisable for these children to have additional immunisations, for example pneumococcal and influenza. This guidance is designed to give general advice to schools and childcare settings. Some vulnerable children may need further precautions to be taken, which should be discussed with the parent or carer in conjunction with their medical team and school health.

Female staff – pregnancy

If a pregnant woman develops a rash or is in direct contact with someone with a potentially infectious rash, this should be investigated by a doctor who can contact the duty room for further advice. The greatest risk to pregnant women from such infections comes from their own child/children, rather than the workplace.

- Chickenpox can affect the pregnancy if a woman has not already had the infection. Report exposure to midwife and GP at any stage of pregnancy. The GP and antenatal carer will arrange a blood test to check for immunity. Shingles is caused by the same virus as chickenpox, so anyone who has not had chickenpox is potentially vulnerable to a case of shingles.
- German measles (rubella). If a pregnant woman comes into contact with german measles she should inform her GP and antenatal carer immediately to ensure investigation. The infection may affect the developing baby if the woman is not immune and is exposed in early pregnancy.
- Slapped cheek disease (fifth disease or parvovirus B19) can occasionally affect an unborn child. If exposed early in pregnancy (before 20 weeks), inform whoever is giving antenatal care as this must be investigated promptly.
- Measles during pregnancy can result in early delivery or even loss of the baby. If a pregnant woman is exposed she should immediately inform whoever is giving antenatal care to ensure investigation.
- All female staff born after 1970 working with young children are advised to ensure they have had two doses of MMR vaccine.

*The above advice also applies to pregnant students.

Immunisations

Immunisation status should always be checked at school entry and at the time of any vaccination. Parents should be encouraged to have their child immunised and any immunisation missed or further catch-up doses organised through the child's GP.

For the most up-to-date immunisation advice and current schedule visit www.publichealth.hscni.net or the school health service can advise on the latest national immunisation schedule.

When to immunise	Diseases vaccine protects against	How it is given
2 months old	Diphtheria, tetanus, pertussis (whooping cough), polio and Hib	One injection
	Pneumococcal infection	One injection
	Rotavirus	Orally
3 months old	Meningococcal B infection	One injection
	Diphtheria, tetanus, pertussis, polio and Hib	One injection
	Rotavirus	Orally
4 months old	Diphtheria, tetanus, pertussis, polio and Hib	One injection
	Pneumococcal infection	One injection
	Meningococcal B infection	One injection
Just after the first birthday	Measles, mumps and rubella	One injection
	Pneumococcal infection	One injection
	Hib and meningococcal C infection	One injection
Every year from 2 years old up to P7	Influenza	Nasal spray or injection
	Diphtheria, tetanus, pertussis and polio	One injection
3 years and 4 months old	Measles, mumps and rubella	One injection
	Cervical cancer caused by human papillomavirus types 16 and 18 and genital warts caused by types 6 and 11	Two injections over six months
14 to 18 years old	Tetanus, diphtheria and polio	One injection
	Meningococcal infection ACWY	One injection

This is the Immunisation Schedule as of July 2016. Children who present with certain risk factors may require additional immunisations. Always consult the most updated version of the "Green Book" for the latest immunisation schedule on www.gov.uk/government/collections/immunisation-against-infectious-disease-the-green-book#the-green-book

From October 2017 children will receive hepatitis B vaccine at 2, 3, and 4 months of age in combination with the diphtheria, tetanus, pertussis, polio and Hib vaccine.

Staff immunisations. All staff should undergo a full occupational health check prior to employment; this includes ensuring they are up to date with immunisations, including two doses of MMR.

Original material was produced by the Health Protection Agency and this version adapted by the Public Health Agency, 12-22 Linenhall Street, Belfast, BT2 8BS. Tel: 0300 555 0114. www.publichealth.hscni.net

Information produced with the assistance of the Royal College of Paediatrics and Child Health and Public Health England.